

A&A manuscript no.
(will be inserted by hand later)

Your thesaurus codes are:
07 ()

ASTRONOMY
AND
ASTROPHYSICS

February 2, 2008

Large frequency drifts during Type I X-ray bursts

Vahid Rezanian

Institute for Advanced Studies in Basic Sciences, Gava Zang, Zanjan 45195, Iran
Department of Physics, University of Alberta, Edmonton AB, Canada T6G 2J1*

Received / accepted

Abstract. We study the spin-down of a neutron star atmosphere during the Type I X-ray burst in low mass X-ray binaries. Using polar cap acceleration models, we show that the resulting stellar “wind” torque on the burning shell due to the flowing charged particles (electrons, protons and ions) from the star’s polar caps may change the shell’s angular momentum during the burst. We conclude that the net change in the angular momentum of the star’s atmosphere can account for rather large frequency drifts observed during Type I X-ray burst.

Key words: stars: neutron – stars: magnetic fields – X-rays: binaries – X-rays: bursts

1. Introduction

It is believed that thermonuclear flashes in pure helium or mixed hydrogen/helium layers on the surface of weakly-magnetic ($B \leq 10^{10}$ G) accreting (at rate $10^{-11} M_{\odot} \text{ yr}^{-1} < \dot{M} < 10^{-8} M_{\odot} \text{ yr}^{-1}$) neutron stars cause the observed Type X-ray bursts in low mass X-ray binaries (LMXBs) (Hansen & Van Horn, 1975; Maraschi & Cavaliere, 1977; Lewin, van Paradijs, & Taam, 1995; Woosley & Taam, 1976; Bildsten, 1998). The accumulated hydrogen and helium on the surface of the neutron star by accretion ignites and burns periodically. General observed features of Type I X-ray bursts, such as burst fluences ($\sim 10^{39}$ erg), burst recurrence times ($\sim$ few hours), and burst durations ($\sim 10 - 100$ s); are successfully explained by spherically symmetric thermonuclear runaway models (Lewin, van Paradijs, & Taam, 1995; Bildsten, 1998). Recent observations from ten LMXBs with the Rossi X-ray Timing Explorer have revealed nearly-coherent oscillations during Type I X-ray bursts with frequencies in the range $\nu_0 \sim 270 - 620$ Hz (van der Klis, 2000; Strohmayer, 2001). Such highly coherent oscillations are interpreted as rotational modulation of surface brightness asymmetries: burning starts locally at some points, creates a brightness

asymmetry, and spreads over the entire stellar surface as the star rotates. This can be understood by noting that the timescales for accretion between bursts is much longer than the burst duration, so the existence of identical conditions over the whole surface of the star for the burning instability to start simultaneously is very unlikely. Therefore, these oscillations provide a direct measurement of the neutron star spin frequency (Strohmayer et al., 1996). Further, observing such oscillations supports the idea that the neutron stars in LMXBs are the progenitors of the millisecond radio pulsars, see Bhattacharya (1995) for a review.

The discovery of such high coherence, large modulation amplitudes, and stable frequency oscillations while providing the convincing evidence that the burst oscillation frequency is related to the neutron star spin frequency (Strohmayer, 2001), however, has brought many new puzzles. An initial puzzle seen in the observations was that the oscillation frequency increases by $\Delta\nu \sim$ a few Hz during the burst. Strohmayer et al. (1997) firstly proposed that this frequency shift is due to the conservation of angular momentum of the decoupled burning shell from the neutron star in which the shell undergoes spin changes as it expands and contracts during the Type I X-ray bursts. Motivated by this proposal, Strohmayer & Markwardt (1999) and Miller (1999, 2000) modeled the observed frequency evolution and found that the oscillations are coherent, as expected if rotation is the cause. But the first theoretical investigations on understanding the problem was done by Cumming & Bildsten (2000) who studied the rotational evolution of the neutron star atmosphere during a thermonuclear burst by considering one-dimensional vertical hydrostatic expansion. Since the nuclear burning time scale ($>$ fraction of a second) is much bigger than the vertical sound crossing time scale through the burning layer ($\sim$ microseconds), the outer layers of the neutron star atmosphere are always in hydrostatic balance. So by assuming conservation of the angular momentum of the shell, a hot burning layer expands hydrostatically by $\Delta R \sim 20$ m (Ayasli & Joss, 1982; Hanawa & Fulimoto, 1984; Bildsten, 1998), and lags behind the neutron star by $\Delta\nu \sim \nu_s(2\Delta R/R) \sim 1\text{Hz}(\nu_s/300\text{Hz})(\Delta R/20\text{m})(10\text{km}/R)$

Send offprint requests to: V. Rezanian

email: vrezania@phys.ualberta.ca

* Present address

where ν_s is neutron star spin frequency and R the radius. As the layer cools down and contracts, the rising oscillations due to a temperature inhomogeneity, drifts upward as we seen by $\Delta\nu$ of few Hertz. By assuming that the burning shell rotates rigidly, they found rough agreement with the observed values with fractional frequency shifts $\Delta\nu/\nu_0 \leq 0.8\%$.

However, recent observations suggested that purely radial hydrostatic expansion and angular momentum conservation alone cannot account for explaining rather large frequency drifts ($\Delta\nu/\nu_0 \sim 1.3\%$) observed in some bursts (Galloway et al., 2001; Wijnands et al., 2001). As pointed out by Galloway et al. (2001), the required expansion by these frequency drifts is 4-5 times larger than the one predicted by Cumming & Bildsten (2000), see Table 1. In order to achieve the rather large frequency shift, Cumming et al. (2001) improved the calculations done by Cumming & Bildsten (2000) by including the general relativistic corrections for either slowly or rapidly rotating star. In agreement with Abramowicz et al. (2001), they found that the general relativity has small effect on the angular momentum conservation law ($\sim 5 - 10\%$). Comparing with the data, for a rigidly rotating atmosphere, they obtained that the expected spin-down is a factor of two or three less than the actual observed values. In another attempt, Spitkovsky et al. (2001) considered a two-dimensional hydrodynamics model that the burning spots can spread over the neutron star surface. Due to the combination of the radial expansion of the burning shell and rotation of the star, they proposed that horizontal hydrodynamics flows may arise in the neutron star burning ocean during the Type I X-ray burst. By taking into account the action of the Coriolis force due to the rapid rotation of the star, they showed that the horizontal flows may explain many features of observed bursts such as the very short rise time of X-ray bursts and the lack of burst oscillations in several bursts. Further, they argued that during cooling the hot ashes left by burning front, there is temperature gradient between equator and pole, increasing toward the pole, which drives a zonal thermal wind directed *backward* to the neutron star rotation.¹ They found that *only if* the frequency drift due to the radial expansion is combined with the geostrophic drift caused by backward zonal flows, one may expect to observe the rather large frequency drifts of burst oscillations in tails of some bursts.

Previous models are based on the angular momentum conservation of the expanding/contracting neutron star atmosphere during the Type I X-ray burst. Although these models are mostly successful in explaining the many features of Type I bursts, the rather large fre-

quency drifts observed in some bursts are still unresolved (Strohmayer & Bildsten, 2003).

Because of no coherent pulsations seen in persistent emission from the majority of neutron stars in LMXBs, it is believed that they are weakly magnetized ($B \leq 10^9$ G). So, in previous studies the effect of magnetic field of the neutron star was ignored in the rotational evolution of burning atmosphere during the X-ray bursts. In this paper we study the change in the angular momentum of the burning shell during the Type I X-ray burst. This study is based on polar cap particle acceleration models in the pulsar polar cap regions. Due to the star's rotation and magnetic field an electric field must be induced in magnetosphere, i.e. $\mathbf{E} = -(\boldsymbol{\Omega}_s \times \mathbf{r}) \times \mathbf{B}/c$, that has non-zero component along the magnetic field lines ($E_{||} = \mathbf{E} \cdot \mathbf{B}$) (see for recent review Mestel, 1998). $\boldsymbol{\Omega}_s$ is the angular velocity of the star, and r is a distance outside of the star. The parallel electric field accelerates charged particles along the open magnetic field lines above the polar caps up to ultrarelativistic energies toward the star's light cylinder ($r_{lc}\Omega = c$). Here, we argue that, for typical magnetic field $B \sim 10^8$ G, the net charged particles flowing from star surface to infinity in the pulsar polar caps, would exert a stellar wind torque on the burning shell that may cause the angular momentum of the shell to change during the burst. We show that the net change in the angular momentum of the shell during the burst can account for rather large frequency drifts of burst oscillations.

2. The model

The theory of particle acceleration in pulsar magnetosphere has been studied for three decades, after pioneering work by Goldreich & Julian (1969) on pulsar electrodynamics. The charged particles accelerated by the parallel electric field (relative to the magnetic field lines) escape from the surface along the open field lines which extended beyond the star's light cylinder (of radius $r_{lc} \sim 5 \times 10^9/\nu_s$ cm, where ν_s in Hertz), and form a relativistic stellar wind. Several kind of acceleration mechanisms has been developed (Mestel, 1998), however, the one proposed by Arons (1979, 1981) may apply likely to X-ray burst pulsars that we will consider here. Arons assumed that charged particles (electrons and ions) flow freely from the neutron star surface due to the thermal activities. Freely escaping particles from the neutron star surface depends on the binding energy (referred as the cohesive energy for ions and as the work function for electrons) and the surface temperature. For typical magnetic field of a neutron star in LMXBs ($\sim 10^8$ G), the threshold temperature for thermionic emission of electrons is $T_e \sim 10^4$ K and for ions is $T_i \sim 10^3$ K (Luo, Shibata & Melrose, 2000) that are $10^4 - 10^5$ lower than surface temperature of the neutron star in LMXBs. Further, due to the thermonuclear activities on the surface of the star during the X-ray burst, the particles' kinetic en-

¹ Actually they showed that due to the hydrostatic balance the material will likely ignite sooner at the equator rather than the pole. As a result, one would expect the burning front cools down at the equator firstly.

ergy would increase significantly. Therefore, freely flowing particles is very likely from the surface of such a pulsar.

Outward flowing charged particles above the pulsar polar caps along the open magnetic field lines causes the Goldreich-Julian (corotation) charge density $\rho_{\text{GJ}} \sim \mathbf{\Omega}_s \cdot \mathbf{B}/2\pi c$ cannot be kept balanced (Harding & Muslimov, 1998). As a result, a strong electric field develops along the magnetic field, $E_{\parallel}$, above the magnetic poles due to the departure of total charge density ρ_e from the Goldreich-Julian charge density, starts from zero at the surface and grows with distance above the surface. Even though $\rho_e = \rho_{\text{GJ}}$, and therefore $E_{\parallel} = 0$ at the surface, the curvature of the field lines causes the area of the open-field region to increase, so that ρ_e increases faster than ρ_{GJ} , and then a charge deficit grows with distance (Harding & Muslimov, 1998). Calculation of the parallel electric field is so complicated and depends on several mechanisms that work to enhance or screen the field. In a series of papers Harding & Muslimov (1998, 2001, 2002); and Harding, Muslimov & Zhang (1998) have extensively studied the influence of pair production, inertial frame-dragging effect, curvature radiation, and inverse Compton scattering on the $E_{\parallel}$. They showed that for the unsaturated region with altitude in range $0 < z \ll (\Omega_s R/c)^{1/2} \simeq 0.1$ (for $\Omega_s = 600\pi(\nu_s/300 \text{ Hz}) \text{ rad/s}$ and $R \sim 10^6 \text{ cm}$) above the stellar surface, the parallel electric field grows linearly with height as $E_{\parallel}(\theta, \phi, z) = E_0(\theta, \phi)z$, while for the saturated region with $z > (\Omega_s R/c)^{1/2}$, the parallel electric field that is nearly constant respect to the altitude, drops by three order of magnitudes. Here θ and ϕ are spherical polar and azimuthal angles, and z is the altitude in units of stellar radius. The amplitude $E_0(\theta, \phi)$ depends on magnetic field strength B , spin frequency ν_s , and orientation of the magnetic field symmetric axis relative to the star spin axis χ . In a simple form, one can estimate $E_{\parallel}$ when the charged particles flow freely from the neutron star surface (see Arons, 1981; Usov & Melrose, 1995; Harding & Muslimov, 2001) as

$$E_{\parallel} \simeq 8.2 \times 10^5 \text{ V/m} \left(\frac{\nu_s}{300 \text{ Hz}} \right)^{5/2} \left(\frac{R}{10^6 \text{ cm}} \right)^{5/2} \times \left(\frac{B}{10^8 \text{ G}} \right) \left(\frac{z}{10^{-4}} \right). \quad (1)$$

Note we calculated $E_{\parallel}$ here for a typical neutron star in LMXBs with nearly perpendicular field orientation relative to the spin axis. The relative fraction of electric force to the gravity force above the burning front during the burst, i.e. $z \sim 10^{-3}$, for electrons and protons are $\sim 8 \times 10^4$ and ~ 42 , respectively. As a result, electrons, protons, and even ionized helium atoms can be accelerated easily along the magnetic field lines by parallel electric field presented in the polar caps of neutron stars in LMXBs. Since the magnetic field lines in the polar cap regions are not closed and extended beyond the star's light cylinder, the accelerated particles in these regions may leave the neutron

star magnetosphere. The resulting flow of particles from the polar caps to infinity produces a relativistic “stellar wind”, that exerts a “stellar wind torque” (Michel, 1991) on the shell, and then causes the net angular momentum of the shell to change during the burst. The net wind torque exerted on the shell due to the out flowing charged particles from the star polar caps is estimated by Michel (1991) for the Goldreich-Julian charge density to be

$$T_W = I^2/4c\Omega_s \quad (2)$$

where $I = \eta n_{\text{GJ}} e v A_{\text{pc}}$ is the total current of the charged particles flowing from the polar caps with the area $A_{\text{pc}} \approx 1.6\pi R^3/R_c$ that is 10% of total star's area ($4\pi R^2$). Here $n_{\text{GJ}} \simeq 0.36\Omega_s B/2\pi e c \simeq 2.25 \times 10^{18}(\nu_s/300 \text{ Hz})(B/10^8 \text{ G}) \text{ cm}^{-3}$ is the Goldreich-Julian charge number density, $e = 1.6 \times 10^{-19} \text{ C}$, $R_c = (GM_s/\Omega_s^2)^{1/3}$ is the corotation distance for a star with mass M_s , and $v \geq 0.1c$ is average speed of particles at $z \sim 10^{-3}$. The velocity of particle grows up quickly to $\sim c$ at $z \sim 0.1$ as it is accelerated by $E_{\parallel}$. $\eta \geq 1$ is a dimensionless factor (free parameter of the model) such that the quantity ηn_{GJ} represents the actual number density of particles that accelerated and left the star due to the parallel electric field. The value of $\eta \simeq 1$ for old and isolated neutron stars (with no observed bursts), i.e. the space charge density is nearly Goldreich-Julian charge density for these stars. But in neutron stars in LMXBs with an ocean of accumulated hot matters, the value of η must be greater than 1, due to the free ejection of particles from the front surface of the ocean. Further, during bursting the accumulated materials with $T \geq 10^8 \text{ K}$ on the neutron star surface, the kinetic energy of particles inside the ocean abruptly increases and then, the space charge density would increase significantly in regions with non-zero parallel electric field. As a result, one would expect that the number particles that may leave the shell would increase during the burst.

It is necessary to note that for neutron stars in LMXBs with accretion rate, $10^{-11} M_{\odot} \text{ yr}^{-1} < \dot{M} < 10^{-8} M_{\odot} \text{ yr}^{-1}$, the existence of pulsar wind torque might not be consistent with accretion. One would expect the motion of the accreting material to oppose and shut off the radio pulsar wind, and contribute to the angular momentum of the system. This scenario is unlikely “during the X-ray burst” due to the following reasons: huge thermonuclear explosion on the surface of the star causes explosive eruption the accumulated materials from the surface during the burst with a mass bulk velocity close to the speed of light. As a result, one would expect the action of accretion to more or less shut off in this period, e.g by evacuation of part of the inner accretion disk during radius expansion. Furthermore, it has been suggested the action of accretion torque is balanced by the gravitational radiation torque in neutron stars in LMXBs, see Bildsten (1998a, 2002) for details. Therefore, the increased angular momentum due to accretion is lost to gravitational radiation. The latter mechanism provides a natural explanation for the rota-

tion of stars in a narrow range of frequencies, $\nu_s \sim 300$ Hz. Finally, the magnitude of accretion torque $T_a \sim \dot{M}(GM_s R)^{1/2} \simeq 8.6 \times 10^{32}$ dyne cm ($\dot{M}/10^{-9} M_\odot \text{yr}^{-1}$) (for $M_s = 1.4 M_\odot$ and $R = 10$ km) is one order of magnitude smaller than that of the wind torque $T_w \sim 9.32 \times 10^{33}$ dyn cm for $\eta \sim 1000$ and $B \sim 10^8$ G, see Eq. (3) and the discussion below Eq. (9).

By replacing I in Eq. (2) we obtain

$$T_w \simeq 9.32 \times 10^{27} \text{ dyne cm } \eta^2 \left(\frac{M_s}{1.4 M_\odot} \right)^{-2/3} \times \left(\frac{\nu_s}{300 \text{ Hz}} \right)^{7/3} \left(\frac{R}{10 \text{ km}} \right)^6 \left(\frac{B}{10^8 \text{ G}} \right)^2. \quad (3)$$

Equation (3) gives the exerted torque on the neutron star surface due to the flowing charged particles from the pulsar polar caps. Therefore, the burning shell of the neutron star will experience this torque as it expands during the burst. The angular momentum of the shell at a distance R from the center of the star in Newtonian dynamics can be written as

$$\ell = \kappa M R^2 \Omega, \quad (4)$$

where constant $\kappa = (I/MR^2)_{\text{shell}}$ depends on the equation of state and $\kappa \leq 2/3$ (an uniform spherical shell). Here M and I are the mass and moment inertia of the shell. Equation (4) may be corrected by $\ell = \kappa M R^2 \sin^2 \theta \Omega_s / (1 - 2GM_s/R)^{1/2}$ for a relativistic spherical star and by $\ell = \kappa M R^2 \sin^2 \theta (\Omega_s - \omega) / (1 - 2GM_s/R)^{1/2}$ for a rotating relativistic star (see Cumming et al., 2001). Here $\omega = J_s/R^3$ where J_s is the total angular momentum of the star. However, as shown by Cumming et al. (2001), the relativistic corrections have small contributions to the final results, and so we ignore them. The time evolution of the angular momentum of the shell will be

$$\frac{\Delta}{\Delta t} (R^2 \Omega)_{\text{shell}} = -T_w / \kappa M, \quad (5)$$

or

$$(R^2 \Omega)_{\text{shell}} (2\Delta R/R + \Delta \Omega/\Omega)_{\text{shell}} = -(T_w / \kappa M) \Delta t. \quad (6)$$

Therefore, the change in the angular velocity of the burning shell will be

$$(\Delta \Omega/\Omega)_{\text{shell}} = -2\Delta R/R - (T_w / \kappa M R^2 \Omega) \Delta t. \quad (7)$$

Here we assumed that the change in the mass of the shell is order of or less than $(\Delta R/R)^2$ and then we neglect it (see Cumming et al., 2001). Equation (7) represent the change in angular velocity of the shell during the burst. Now we discuss the change in angular velocity in three stages, before the X-ray burst or the recurrence period, the rising period, and after the burst.

2.1. Before the X-ray burst

In this period the star accumulates materials in the ocean that more or less rigidly corotates with the star (Cumming et al., 2001). The parallel electric field inside the ocean is nearly neutralized due to redistribution of charge particles of the ocean. So in this period, the electric field more or less start from zero above the front surface of the shell and increase by height due to the charge deficit caused by free emission of charged particles from the front surface. The space charge density above the front surface would be bigger than but not too far from the Goldreich-Julian charge density and increases by time due to the thermonuclear activities in the oceans. So we expect that $\eta \sim 10 - 100$.

However, the polar cap acceleration is more or less suppressed due to strongly accreting plasma in this period. As a result, the pulsar wind torque would be negligible.

Furthermore, because of nearly rigid rotation of the ocean with star the net wind torque during this period will be acted more or less on the star itself. Therefore, Eq. (7) should be corrected for this period as

$$\begin{aligned} \left(\frac{\Delta \Omega}{\Omega} \right)_{\text{shell}} &= -(T_w / \kappa' M_s R^2 \Omega) \Delta t, \\ &\simeq -4.44 \times 10^{-17} \left(\frac{\eta}{100} \right)^2 \left(\frac{M_s}{1.4 M_\odot} \right)^{-5/3} \\ &\times \left(\frac{R}{10 \text{ km}} \right)^4 \left(\frac{\nu_s}{300 \text{ Hz}} \right)^{4/3} \left(\frac{B}{10^7 \text{ G}} \right)^2 \left(\frac{\Delta t_{\text{rec}}}{10^4 \text{ s}} \right), \end{aligned} \quad (8)$$

where Δt_{rec} is the burst recurrence time and $\kappa' = 2/5$. As is clear, the net wind torque on the shell/star in this period and so the change in angular velocity of the shell/star is negligible.

2.2. The rising time

This period starts with abruptly thermonuclear ignition of the accumulated hydrogens and helium during the recurrence time. The burning layers expand in less than ~ 1 second up to $z \simeq 2 \times 10^{-3}$ (Ayasli & Joss, 1982; Hanawa & Fulimoto, 1984; Bildsten, 1998). In this period, as in the hydrostatic models, we assume that the rotational evolution of the expanding shell is nearly independent of the star itself.

As we mentioned before, in this period due to the huge thermonuclear explosion on the surface of the star, the accumulated materials are explosively erupted out of the star with a mass bulk velocity close to the speed of light. The outgoing plasma density within period of time $\Delta t \sim 1$ second, is $\sim [(A_{\text{pc}}/4\pi R^2) M_{\text{shell}}]/A_{\text{pc}}$ that is 10 times (or more) larger than in-falling plasma density $m_{\text{acc}}/A_{\text{pc}}$ due to the accretion flow at the same period of time. Here $m_{\text{acc}} = \dot{M} \Delta t$ is the average mass of accreting materials during Δt . As a result, one would expect the action of accretion to more or less shut off during this period.

The parallel electric force at height $z \simeq 2 \times 10^{-3}$ is 80 times bigger than gravitational force for hydrogen atoms, see Eq. (1). Therefore, the charged particles that rise with the expanding layers will be accelerated toward the outside of light cylinder by the strong parallel electric field along the open magnetic field lines. Further, the sudden thermonuclear explosion of the accumulated materials in the ocean causes the thermal energy of the charged particles to increase. Consequently, we expect that the number density of particles that may leave the magnetosphere ηn_{GJ} in this period increases significantly. The resulting wind torque for $\eta \sim 10^3$ will give a net change in angular velocity of the shell as

$$\begin{aligned} \left(\frac{\Delta\Omega}{\Omega} \right)_{\text{shell}} &= -2|\Delta R|/R - (T_w/\kappa M R^2 \Omega) \Delta t, \\ &\simeq -2 \times 10^{-3} \left(\frac{z}{10^{-3}} \right) \\ &\quad - 7.42 \times 10^{-3} \left(\frac{\eta}{10^3} \right)^2 \left(\frac{M_s}{1.4 M_\odot} \right)^{-2/3} \left(\frac{M}{10^{21} \text{ g}} \right)^{-1} \\ &\quad \times \left(\frac{R}{10 \text{ km}} \right)^4 \left(\frac{\nu_s}{300 \text{ Hz}} \right)^{4/3} \left(\frac{B}{10^8 \text{ G}} \right)^2 \left(\frac{\Delta t_{\text{rise}}}{1 \text{ s}} \right), \end{aligned} \quad (9)$$

where $\kappa = 2/3$ and $\Delta R \simeq 10 \text{ m}$ is the average change in thickness of the burning layers during a burst. For $\eta \sim 10^3$ and $B \sim 10^8 \text{ G}$, the net change in angular velocity of the shell given by Eq. (9) is in good agreement with large frequency shifts seen in observations. We note that for $\eta \sim 10^3$, the number of particles that left the star $\Delta N \sim 4\pi\eta n_{\text{GJ}} R^2 \Delta R$ is much smaller than total number of particles in the ocean $N \sim M/m_p$, i.e. $\Delta N/N \simeq 4 \times 10^{-7}$. As expected, the change in mass of the shell is order of $(\Delta R/R)^2 \sim 10^{-6}$. In Table 1 we calculate the quantity ηB for both 300 Hz and 600 Hz spin frequencies (for those which applicable), to produce the frequency shifts observed in various systems. As a result, the suggested magnetic fields are in range $1 - 10 \times 10^7 \text{ G}$ for $\eta \sim 100 - 1000$. This is consistent with other observational evidences about the magnetic field of neutron stars in LMXBs.

2.3. After the X-ray burst

At final stage, after the thermonuclear flash, the expansion will be stopped by the star's gravity force, and then the expanded layers start to contract as they cool down. Because of the charge redistributions during the expansion, the original induced electric field is neutralized inside the expanded shell. So the magnitude of the parallel electric field inside the ocean will drop to zero, and then the electrostatic acceleration will cease inside the ocean. Due to the lack of outward electrostatic acceleration, the ocean's particles are mostly accelerated downward by strong gravity. Furthermore, the action of accretion plasma is starting to recover, and as a result, the polar cap acceleration will shut down in this period. Although, because of new

charge distribution, a new parallel electric field may develop above the burned front we expect that the polar cap particle acceleration mechanism is more or less unlikely after the burst. As a result, the number of particles that may leave the star will decrease as well, i.e. $\eta \leq 10$. Further as the shell cools down, it gets closer and closer to the star, and then its coupling to the star gets stronger and stronger. This can be understood by noting the magnetic field near the star's surface opposes the differential rotation arising between shell's layers and surface of the star during the burst. Therefore, the net wind torque will be acting more and more on the whole star again rather than shell itself, and so we neglect it, see Sect. 2.1.

The magnetic recoupling of the shell forces the shell to spin up until it achieves the star's spin frequency. The shell gains its angular momentum deficit $(MR^2\Delta\Omega)_{\text{shell}}$, from the star, and then it reduces the angular velocity of the star by $\delta\Omega_s = (M/M_s)\Delta\Omega \simeq 10^{-12}\Delta\Omega$ which is negligible. As a result, one would expect that the oscillations' frequency will increase as the shell spins up by $\Delta\nu/\nu_0 \sim \Delta\Omega/\Omega$. In Fig. 1, we compare the corresponding changes suggested by previous studies (Strohmayer et al., 1997; Cumming et al., 2001) and the one we discussed here. As is clear, due to the exerted torque by the particle wind in the polar cap regions, the shell's spin frequency, and then the oscillations' frequency decrease more than in the calculation by Cumming et al. (2001) during the rising time. Therefore, we expect to observe larger change in oscillations' frequency during the burst tail.

3. Discussion

Among the ~ 50 known Galactic LMXBs, the highly coherent burst oscillations with large modulation amplitudes and stable frequencies in range $\nu_0 \sim 270 - 620 \text{ Hz}$ are seen in ten LMXBs during Type I X-ray bursts. These oscillations are most commonly seen during tails of bursts, when the burning is thought to have spread over the whole surface and obviously asymmetry is no longer present, are not observed in all Type I X-ray bursts from the same source. While it is believed that the oscillation frequency is closely related to the neutron star spin frequency due to the observation of the kHz quasi-periodic oscillations in the persistent emission (van der Klis, 2000), there is still an ambiguity that whether the oscillation frequency is the spin frequency or twice the spin frequency. Further, as seen in observations, the oscillation frequency increases by $\Delta\nu$ a few Hz during the burst. This frequency shift is firstly explained by Strohmayer et al. (1997) that the burning shell decouples from the star, and undergoes spin changes due to the conservation of angular momentum of the shell as it expands and contracts during the Type I X-ray bursts. Further observations and studies suggested that purely radial hydrostatic expansion and angular momentum conservation alone cannot explain rather large frequency drifts ($\Delta\nu/\nu \sim 1.3\%$) observed in some bursts (Cumming et al.,

2001; Galloway et al., 2001; Wijnands et al., 2001). For recent review on the thermonuclear bursts and their properties see Strohmayer & Bildsten (2003).

In this paper, we addressed the latter problem by studying the evolution of angular momentum of the burning shell during the Type I X-ray bursts in LMXBs. Based on particle acceleration models near a pulsar polar cap region (Mestel, 1998), we studied the change in the angular momentum of the burning shell during the Type I X-ray burst. The net charged particles that accelerated by parallel electric field in star's polar caps, flow from the star surface to infinity through open magnetic field lines, would exert a torque on the burning shell (called wind torque) and cause the angular momentum of the shell changes during the burst. We introduced a dimensionless factor η that the quantity ηn_{GJ} represent the number density of the ejected particles from the burning layers above the pulsar polar caps. We showed that for $\eta \sim 100 - 1000$ (during X-ray bursts) and a typical magnetic field $B \sim 1 - 10 \times 10^7$ G in the pulsar polar caps, the rather large observed frequency drifts of burst oscillations can be explained by the resulting wind torque exerted on the burning shell. In Table 1, we obtained the values for ηB for both 300 Hz and 600 Hz spin frequencies that causes the corresponding frequency drift observed in each burst. As is clear, the resulting magnetic fields' strength is in a good agreement with other observations from the neutron stars in LMXBs, such as the lack of coherence pulsations in persistent emission.

Above the neutron star's polar caps particles flow outward along the open magnetic field lines and a steady charge density cannot be maintained at the Goldreich-Julian density everywhere. Consequently, a strong electric field develops along the magnetic field, extracts the 'primary' charged particles with density $n \sim n_{\text{GJ}}$ (for a star with no burst activity, see Sect. 2) from the surface and accelerate them to high Lorentz factor ($\sim 10^7$). According to the polar cap models, the predicted radiation luminosity for a typical millisecond pulsar would be $\mathcal{E} \sim 10^{33}(\nu_s/300 \text{ Hz})^{5/28}(B/10^8 \text{ G})^{-1/7} \text{ ergs/s}$ that is comparable to γ -ray luminosities. So, one may expect to observe γ -rays in millisecond pulsars that has been detected only in PSR J0218+4232 by EGRET (Kui et. al., 2000). However, polar cap particle acceleration is limited by various energy loss mechanisms such as curvature radiation, resonant and nonresonant inverse Compton scattering, and pair production. The accelerated particles along the magnetic field lines suffer energy loss and emit photons through the curvature radiation. Further, these particles scattered by soft thermal X-ray photons emitted from the hot surface and loss energy via inverse Compton scattering. In addition, the high energy photons resulted from the curvature radiation and/or inverse Compton scattering produce a cascade of 'secondary' $e^\pm$ pair particles with a number density $n_{e^\pm} \sim 10^3 n$ above polar cap region that change the charge distributions and screen the electric field (see for more detail Harding & Muslimov,

2001). In millisecond pulsars (with no burst activities) the polar cap particle acceleration mechanism was studied by Luo, Shibata & Melrose (2000). They showed that in these pulsars this mechanism, which is relatively efficient in accelerating charged particles in the star's polar caps, mostly suffers energy loss from the radiation of accelerated particles along the field lines (curvature radiation). So the maximum energy attainable by the particle is limited by energy loss through the radiation reaction. They found that the maximum Lorentz factor that might be achieved by an individual particle at $z \sim 0.1$ is $\Gamma = (6\pi\epsilon_0 E_{\parallel} \rho_c^2/e)^{1/4} \sim 10^7$, however, the Lorentz factor for the bulk pair plasma is much smaller ~ 100 . The latter may explain the lack of detection of high energy γ -rays in these stars. Here $E_{\parallel}$ is given by Eq. (1) and $\rho_c \sim (4/3)(cR/2\pi\nu_s)^{1/2} \sim 5 \times 10^4(300 \text{ Hz}/\nu_s)^{1/2}(R/10^6 \text{ cm})$ m is the curvature radius of the field lines.

For neutron stars in LMXBs with intermediate accretion rate $10^{-11}M_{\odot} \text{ yr}^{-1} < \dot{M} < 10^{-8}M_{\odot} \text{ yr}^{-1}$, however, the polar cap acceleration mechanism mostly suffers from the action of accretion plasma. Because of the large plasma density ($m_{\text{acc}}/A_{\text{pc}}$) of the accretion flow toward star's polar caps, the parallel electric field in the polar cap regions would be shorted out and the polar cap acceleration mechanism would be suppressed. We note that, this scenario is not likely during X-ray bursts due to the explosively erupting materials from the surface and possible evacuation of part of inner disk. In this period with $\Delta t \sim 1$ second, the eruptive plasma density is 10 times larger than the accreting plasma density, i.e. $\sim [(A_{\text{pc}}/4\pi R^2)M_{\text{shell}}]/A_{\text{pc}} \sim 10 m_{\text{acc}}/A_{\text{pc}}$.

Acknowledgements. The author is grateful to Sharon M. Morsink and Supratim Sengupta for reading the manuscript and stimulating discussions during the course of this work. I would like also thank to the referee for his/her valuable comments. This research was supported by the Natural Sciences and Engineering Research Council of Canada.

References

- Abramowicz, M. A., Kluzniak, W., & Lasota, J.-P. 2001, *A&A*, 374, L16
- Arons, J. 1979, *Space Sci. Rev.*, 24, 437
- Arons, J. 1981, *ApJ*, 248, 1099
- Ayasli, S., & Joss, P. C. 1982, *ApJ*, 256, 637
- Bhattacharya, D. 1995, in *X-ray Binaries*, ed. W. H. G. Lewin, J. van paradijs, & E. P. J. van den Heuvel (Cambridge: Cambridge University Press), 233
- Bildsten, L. 1998, in *The Many Faces of Neutron Stars*, ed. R. Bucccheri, J. van Paradijs, & M. A. Alpar (Dodrecht: Kluwer), 419
- Bildsten, L. 1998, *ApJ*, 501, L89
- Bildsten, L. 2002, to appear in "Radio Pulsars" (ASP Conf. Ser.), eds M. Bailes, D.J. Nice, & S.E.Thorsett; pre-print: *astroph/0212004*
- Cumming, A. & Bildsten, L. 2001, *ApJ*, 559, 127
- Cumming, A., Morsink, S. M., Bildsten, L., Friedman, J. L., & Holz, D. E. 2002, *ApJ*, 564, 343

Fox, D. W., Muno, Lewin, W. H. G., M. P., Morgan, E. H., & Bildsten, L. 2001, AAS, 198, 270

Galloway, D. K., Chakrabarty, D., Muno, M. P., & Savov, P. 2001, ApJ, 549, L85

Goldreich, P., & Julian, W. H. 1969, ApJ, 157, 869

Hanawa, T. & Fujimoto, M. Y. 1984, PASJ, 36, 199

Hansen, C. J. & Van Horn, H. M. 1975, ApJ, 317, 368

Harding, A. K. & Muslimov, A. G. 1998, ApJ, 508, 346

Harding, A. K. & Muslimov, A. G. 2001, ApJ, 556, 987

Harding, A. K. & Muslimov, A. G. 2002, ApJ, 568, 877

Harding, A. K., Muslimov, A. G., & Zhang, B. 2002, ApJ, 576, 366

Kuiper, L., et. al. 2000, A&A, 359, 615

Lewin, W. H. G., van Paradijs, J., & Taam, R. E. 1995, in X-ray Binaries, ed. W. H. G. Lewin, J. van paradijs, & E. P. J. van den Heuvel (Cambridge: Cambridge University Press), 175

Luo, Q., Shibata, S., & Melrose, D. B. 2000, MNRAS, 318, 943

Maraschi, L. & Cavaliere, A. 1977, Highlights Astron., 4, 127

Mestel, L. 1998, Stellar Magnetism (Oxford: Oxford Univ. Press)

Michel, F. C. 1991, Theory of Neutron Star Magnetosphere, (Chicago: The University of Chicago Press)

Miller, M. C. 2000, ApJ, 515, L77

Miller, M. C. 2000, ApJ, 531, 458

Muno, M. P., Chakrabarty, D., Galloway, D. K., & Savov, P. 2001, ApJ, 553, L157

Muno, M. P., Fox, D. W., Morgan, E. H., & Bildsten, L. 2000, 542, 1016

Smith, D. A., Morgan, E. H., & Bradt, H. 1997, ApJ, 479, L137

Spitkovsky, A., Levin, Y., & Ushomirsky, G. 2002, ApJ, 566, 1018

Strohmayer, T. E. 2001, Advances in Space Research, Vol. 28, Issue 2-3, 511, preprint:astro-ph/0012516

Strohmayer, T. E. 1999, ApJ, 523, L51

Strohmayer, T. E., & Bildsten, L. 2003, preprint:astro-ph/0301544

Strohmayer, T. E., Zhang, W., Swank, J. H., & Lapidus, I. 1998, ApJ, 498, L135

Strohmayer, T. E., Jahoda, K., Giles, B. A., & Lee, U. 1997, ApJ, 486, 355

Strohmayer, T. E., Zhang, W., Swank, J. H., Smale, A., Titarchck, L., Day, C., & Lee, U. 1996, ApJ, 469, L9

Strohmayer, T. E., & Markwardt, C. B. 1999, ApJ, 516, L81

Usov, V. V., & Melrose, D. B. 1995, preprint: astro-ph/9506021

van der Klis, M. 2000, ARA&A, 38, 717

van Straaten, S., van der Klis, M., Kuulkers, E., & Mendez, M. 2001, ApJ, 551, 907

Wijnands, R., Strohmayer, T. E., & Franco, L. 2001, ApJ, 549, L71

Woosley, S. E. & Taam, R. E. 1976, Nature, 263, 101

Zhang, W., Jahoda, K., Kelley, R. L., Strohmayer, T. E., Swank, J. H., & Zhang, S. N. 1998, ApJ, 495, L9

– Table 1: In this table we obtain the lowest radius expansion ΔR (based on hydrostatic expansion models) and the scaled magnetic field ηB (based on the polar cap particle acceleration models) for some X-ray bursts in LMXBs. We calculate these quantities for two chosen values of the star spin frequency, $\nu_s = 300$ Hz and $\nu_s = 600$ Hz, such that the corresponding frequency shift is comparable with observations. ν_0 is the oscillation frequency and $\Delta\nu$ its corresponding shift seen during the burst. The value of ηB is based on the assumed value of $\Delta R \simeq 20$ m for both spin frequencies, see Eq. (9).

Table 1

Object	Time (UT)	ν_0 (Hz)	$\Delta\nu/\nu_0$ (10^{-3})
4U 1636-54	1996 Dec 28 (22:39:22) ^{1,2}	580.5	$\sim 2-4$
	1996 Dec 29 (23:26:46) ^{2,3}	581.5	~ 3
	1996 Dec 31 (17:36:52) ^{2,3}	581.	~ 3
4U 1702-43	1997 Jul 26 (14:04:19) ⁴	329.85 ± 0.1	7.7 ± 0.3
	1997 Jul 30 (12:11:58) ⁴	330.55 ± 0.02	4.8 ± 0.3
4U 1728-34	1996 Feb 16 (10:00:45) ^{4,5}	364.23 ± 0.05	6.6 ± 0.1
	1997 Sep 9 (06:42:56) ⁴	364.10 ± 0.05	5.9 ± 0.2
Aql X-1	1997 Mar 1 (23:27:39) ^{6,7}	549.76 ± 0.04	4.3
4U 1916-053	1998 Aug 1 (18:23:45) ⁸	269-272	13.0
MXB 1658-298	1999 Apr 14 (11:44:52) ⁹	567	9.0

References: (1) Strohmayer et al. (1998); (2) Miller (2000); (3) Strohmayer (1999); (4) Strohmayer & Markwardt (1999); (5) Strohmayer et al. (1996); (6) Zhang et al. (1998); (7) Fox et al. (2000); (8) Galloway et al. (2001); (9) Wijnands et al. (2001)




Fig. 1. In this figure we schematically compare the change in angular velocity of the expanding shell and then oscillations' frequency during the type I X-ray burst suggested by Cumming et al. (2001) (dotted line) and the one proposed here (solid line). Due to the exerted torque on the shell by the particle wind from the magnetic polar cap regions during the burst, we expect a larger depth in $\Delta\Omega$ rather than the model discussed by Cumming et al. (2001). As a result, the larger drift would be expected in the burst tail. The solid and dotted curve after the burst are drawn schematically to show star-shell recoupling in this period.